

## References:

- Baker, C., Worrall, L., Rose, M., Hudson, K., Ryan, B., & O'Byrne, L. (2017). A systematic review of rehabilitation interventions to prevent and treat depression in post-stroke aphasia. *Disability and Rehabilitation, 40*(16), 1870–1892. <https://doi.org/10.1080/09638288.2017.1315181>
- Baker, C., Worrall, L., Rose, M., & Ryan, B. (2019). Stroke health professionals' management of depression after post-stroke aphasia: A qualitative study. *Disability and Rehabilitation, 43*(2), 217–228. <https://doi.org/10.1080/09638288.2019.1621394>
- Benaim, C., Decavel, P., Bentabet, M., Froger, J., Péliissier, J., & Pérennou, D. (2010). Sensitivity to change of two depression rating scales for stroke patients. *Clinical Rehabilitation, 24*(3), 251–257. <https://doi.org/10.1177/0269215509346087>
- Code, C., Hemsley, G., & Herrmann, M. (1999). The emotional impact of Aphasia. *Seminars in Speech and Language, 20*(01), 19–31. <https://doi.org/10.1055/s-2008-1064006>
- Griffin-Musick, J. R., Off, C. A., Milman, L., Kincheloe, H., & Kozlowski, A. (2020). The impact of a university-based intensive comprehensive aphasia program (ICAP) on psychosocial well-being in stroke survivors with aphasia. *Aphasiology, 35*(10), 1363–1389. <https://doi.org/10.1080/02687038.2020.1814949>
- Laures-Gore, J. S., Farina, M., Moore, E., & Russell, S. (2016). Stress and depression scales in aphasia: Relation between the Aphasia Depression Rating Scale, stroke Aphasia depression questionnaire-10, and the perceived stress scale. *Topics in Stroke Rehabilitation, 24*(2), 114–118. <https://doi.org/10.1080/10749357.2016.1198528>
- Robinson, R. G., & Jorge, R. E. (2016). Post-stroke depression: a review. *American Journal of Psychiatry, 173*(3), 221-231.
- Schmid, A. A., Kroenke, K., Hendrie, H. C., Bakas, T., Sutherland, J. M., & Williams, L. S. (2011). Poststroke depression and treatment effects on functional outcomes. *Neurology, 76*(11), 1000–1005. <https://doi.org/10.1212/WNL.0b013e318210435e>
- Shehata, G. A., El Mistikawi, T., Risha, A. S., & Hassan, H. S. (2015). The effect of aphasia upon personality traits, depression and anxiety among stroke patients. *Journal of Affective Disorders, 172*, 312–314. <https://doi.org/10.1016/j.jad.2014.10.027>
- Tiwari, V., Chaturvedi, P., Singh, A. K., Qavi, A., & Thacker, A. K. (2020). Depression impedes neuroplasticity and quality of life after stroke. *Journal of Family Medicine and Primary Care, 9*(8), 4039. [https://doi.org/10.4103/jfmpe.jfmpe\\_273\\_20](https://doi.org/10.4103/jfmpe.jfmpe_273_20)

- Towfighi, A., Ovbiagele, B., El Husseini, N., Hackett, M. L., Jorge, R. E., Kissela, B. M., Mitchell, P. H., Skolarus, L. E., Whooley, M. A., & Williams, L. S. (2017). Poststroke depression: A scientific statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*, *48*(2).  
<https://doi.org/10.1161/str.0000000000000113>
- Vasiliu, O., Marinescu, I., & Vasile, D. (2019). Pharmacological therapy of post-stroke depression. *Psihiatru.ro*, *2*(57), 18. <https://doi.org/10.26416/psih.57.2.2019.2441>
- Wijeratne, T., & Sales, C. (2021). Understanding why post-stroke depression may be the norm rather than the exception: The anatomical and neuroinflammatory correlates of post-stroke depression. *Journal of Clinical Medicine*, *10*(8), 1674.  
<https://doi.org/10.3390/jcm10081674>
- Zanella, C., Laures-Gore, J., Dotson, V. M., & Belagaje, S. R. (2022). Incidence of post-stroke depression symptoms and potential risk factors in adults with aphasia in a comprehensive stroke center. *Topics in Stroke Rehabilitation*, 1–11.  
<https://doi.org/10.1080/10749357.2022.2070363>